

Investigator: A. Bishaya

Granulocyte Macrophage Colony Forming Unit (GM-CFU) Assay

Experiment # 8 (^{CB-151} Survival #1) Source of Irradiation: External
 Mice Sex, Strain, Age: SP, F56 wP₁ Irradiation
 Type of Irradiation: Chronic External
 Animals per group: 3

Aims: To determine the bone marrow GM-CFU response to chronic external ¹³⁷Cs irradiation (dose rate ~~of~~ reduction half life = 223h)

Summary of Results:

Corresponds to Sn-117m Tc in femur

Brief Procedure:

- 1) Irradiate animals in groups of 3 with desired initial dose rates (cGy/h) listed in Table 1.
- 2) Sacrifice each mouse on optimal day by cervical dislocation and sterilize using 70% EtOH and immediately move it into laminar flow hood.
- 3) Remove both femurs carefully using sterile instruments and clean the attached tissue thoroughly.
- 4) Flush the bone marrow with 2% Horse Serum in Dulbecco's Modified Eagles Medium (2% HS-DMEM) using 21G needle and syringe.
- 5) Separate the mononuclear cells by density gradient procedures using Histopaque.
- 6) Plate the desired number of in mixture of 60% HS-DMEM and 0.6% bacto agar solution in the presence of 9.2 U (New Sigma unit) GM-CFS.
- 7) Keep the plated well plates for 20 min. in laminar flow hood and move them into incubator with 5% CO₂ and 95% air, at 37°C.
- 8) Count the granulocyte macrophage colonies on 7th day.

Group # Probe #	Starting date	Initial Dose Rate (R/h)	Date Sacrificed	# of days	Remarks if any
C1	09/29/98	0	10/6/98	7	} unirradiated control
C2	09/29/98	0	10/6/98	7	
C3					
1	09/29/98	3.0	10/6/98	7	Cage # 1
2	09/29/98	1.506	10/6/98	7	Cage # 2
3	09/29/98	0.825	10/6/98	7	Cage # 3
4	09/29/98	0.456	10/6/98	7	Cage # 4
5					
6					
7					

Preparing Media and Agar:

Culture Medium (Double Strength): 13.37g (1 pack) of D-MEM powder (Gibco, Cat # 12100-046) + 490 ml deionized water + 16 μ l of L-asparagine (Gibco Cat # 12416-012) at a concentration of 5 μ g/ μ l + 150.4 μ l of DEAE dextran (mol. wt. = 2×10^6 , intrinsic viscosity = 0.7) at a concentration of 1 μ g/ μ l (Sigma Cat # D-9885) + 10 ml of penstrep (Gibco Cat # 600-5070, 5,000 units/ml pen, 5,000 μ g/ml streptomycin) + 3.7 g of NaHCO₃ (Gibco Cat # 11810-025).

Wash Medium: i) Mix equal amounts of culture medium and sterile deionized water.
ii) Add 2% HS

Agar: Prepare 0.6% agar by adding 0.6 g Difco Bacto agar (Difco Cat # 0140-15-4) to 100 ml boiling deionized water. Autoclave on liquid cycle for 20 min.

Comments If any:**Flushing Bone marrow:**

- 1) Remove both femurs from each mouse and place them in a test tube containing wash medium kept in ice, if the femur can not be flushed immediately.
- 2) Flush the marrow from each femur by aspirating 3 ml of Wash Medium through the femur 5 times with a 21G needle/3 ml syringe in a 50 ml conical centrifuge tube. Follow with two flushes with 1 ml of fresh Wash Medium.
- 3) Spin the cells at 1200 rpm for 5 minutes at 4°C, decant, break up the cell pellet, resuspend the cells in 5 ml of cold Wash Medium, and vortex the cell suspension.

Comments If any:**Separating Mononuclear cells and washing the cells:**

- 1) Transfer 3.5 ml of Histopaque (Sigma Cat #H8889) into fresh 15 ml tubes (1 tube per group).
- 2) Layer the cell suspension carefully on top of Histopaque and centrifuge at 1500 rpm, 4°C, for 30 minutes.
- 3) Using a Pasteur pipette transfer the mononuclear cells into fresh tubes.
- 4) Dilute the cell suspension to 15 ml by adding cold Wash Medium into each tube and spin them at 1200 rpm, 4°C, for 5 min.
- 5) Decant the supernatant, break the pellet, and add 15 ml cold Wash Medium, and spin them again at 1200 rpm, 4°C, for 5 min. Repeat this procedure 2 more times.
- 6) After 3rd wash break the pellet and resuspend in 2 ml Culture medium (2x DMEM) with 60% HS and keep the tubes in dry bath at 37°C.
- 7) Add 20 μ l of cell suspension to 20 ml of Isotone II in a coulter cup and determine total # of cells in each group using coulter counter.

Coulter Counter Parameters:Current(I)=500 μ A

Full Scale = 1

T_L = 2.7T_u = 99.9

Attenuation= 4

Alarm Threshold = off

Preset Gain = 1

Stirrer control = off

Coulter Counter Parameters: *Same as above*

ml 2000

Multiplication Factor to get # of cells/ μ l = ~~2~~ Coulter Count

Group #	Coulter Count without ZG	Avg	# cells per μ l	Coulter Count with 5 drops ZG	Avg	# cells per μ l <i>ml</i>
C1				6510, 6439, 6552		} 14,555 <i>000</i>
C2				8013, 8060, 8091		
C3						
1				3379, 3183, 3272	3278	6556000
2				4480, 4372, 4497	4449	8899333
3				3423, 3524, 3374	3440	6880666
4				5222, 5024, 5003	5083	10166000
5						
6						
7						

Average # of cells per μl = $14,555,000$ cells

DILUTIONS

Dilution A: (1.0×10^6 cells / ml, Total volume 3.4 ml)

1.7 ml Agar + 1.47 μl Medium + 0.233 μl Cell Suspension

Dilution B: (3.0×10^5 cells / ml, Total volume 3.4 ml)

1.7 ml Agar + 1.63 μl Medium + 0.07 μl Cell Suspension

~~Dilution C: (1.0×10^5 cells / ml, Total volume 3.4 ml)~~

~~1.7 ml Agar + μl Medium + μl Cell Suspension~~

Plating the Cells:

Culture Medium: Maintain four 13mm tubes each containing 4.5 ml of Culture medium in dry bath at 37°C .

Horse Serum: Maintain five 13mm tubes each containing 4.5 ml of Horse Serum in dry bath at 37°C .

Agar: Maintain five 16mm tubes each containing 6.5 ml of Agar in dry bath at 37°C .

- 1) Warm up dilution tubes (one per group) to 37°C in dry bath.
- 2) Warm up Agar (30 ml) and 60% HS in 2x DMEM (30 ml) in separate tubes to 37°C .
- 3) Mark the well plates (3 wells for each dilution for each group)
- 4) Mix 1.7ml agar + x ml of 2x DMEM with 60% HS + y ml cell suspension + 0.02 ml GMCSF ($x + y = 1.7$ ml) in a dilution tube.
- 5) Add 1 ml of mixture 4 to each well, mix properly and let it gel for about 15 minutes.
- 6) Repeat steps 4 and 5 for each dilution.
- 7) Repeat steps 1 to 6 for each group.
- 8) Incubate the cells in an incubator at 37°C and 5% CO_2 , 95% air for 7 days.
- 9) On 8th day of incubation count colonies and determine the survival fractions.

Comments If any:

Counting the Colonies: (Inverted at 40X or dissecting at 35X)

Group #	Dose (p) Activity, # injected	# of cells plated	# CFU-GM counted	Avg	SF
C1	0	3×10^5	90, 83, 92	95.16	
C2	0	3×10^5	111, 89, 106		
C3					
1	383	1×10^6	30, 32, 27	8.9	0.0944
2	192	3×10^5	21, 18, 21	20	0.2101
3	105	3×10^5	36, 40, 36	37.33	0.3922
4	58	3×10^5	64, 64, 58	62	0.6514
5					
6					
7					